

XXIV PHYSICS IN COLLISION - Boston, June 27-29, 2004

B_d MIXING AND PROSPECTS FOR B_s MIXING AT DØ

Tulika Bose

Columbia University, New York, New York, NY 10027, USA
for the DØ Collaboration

ABSTRACT

Measurement of the B_s oscillation frequency via B_s mixing analyses provides a powerful constraint on the CKM matrix elements. The study of B_d oscillations is an important step towards B_s mixing and a preliminary measurement of Δm_d has been made with $\sim 250 \text{ pb}^{-1}$ of data collected with the upgraded Run II DØ detector. Different flavor tagging algorithms have been developed and are being optimized for use on a large set of B_s mesons that have been reconstructed in different semileptonic decay modes.

1 Introduction

First observed in the $B_d^0 - \bar{B}_d^0$ meson system at ARGUS, the neutral B meson transition from the particle to anti-particle state, and vice versa, occurs through a second order weak transition or “box diagram”. The frequency of the oscillation is proportional to the small difference in mass between the two eigenstates, Δm , and for the $B_d^0 - \bar{B}_d^0$ system can be translated into a measurement of the CKM element $|V_{td}|$. $|V_{td}|$ can be used to constrain the unitarity triangle and thereby yield information on the CP violating phase [1]. Δm_d has been precisely measured (the world average is $\Delta m_d = 0.502 \pm 0.007 \text{ ps}^{-1}$ [2]) but large theoretical uncertainties dominate the extraction of $|V_{td}|$ from Δm_d . This problem can be reduced if the

B_s^0 mass difference, Δm_s , is also measured. $|V_{td}|$ can then be extracted with better precision from the ratio:

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m(B_s^0)}{m(B_d^0)} \xi^2 \left| \frac{V_{ts}}{V_{td}} \right|^2 \quad (1)$$

where ξ is estimated from Lattice QCD calculations to be $1.15^{+0.12}_{-0.00}$ [3]. The above has motivated many experiments to search for B_s^0 oscillations and though a statistically significant signal hasn't been observed yet, a lower limit ($\Delta m_s > 14.4 \text{ ps}^{-1}$ at 95% C.L.) has been set. Since this current limit indicates that the B_s^0 oscillations are at least 30 times faster than the B_d^0 oscillations, a B_s^0 mixing measurement is experimentally very challenging.

2 Experimental considerations

The DØ experiment at the Fermilab Tevatron, a $p\bar{p}$ collider at 1.96 TeV center of mass energy, is well equipped to search for B_s^0 oscillations. The large muon acceptance and forward tracking coverage of the DØ detector (pseudorapidity coverage of $|\eta| < 2.0$ for the muon, $|\eta| < 1.7$ for the tracking and $|\eta| < 3.0$ for the silicon sub-detectors), along with a robust muon trigger are highly effective in exploiting the large $b\bar{b}$ cross-section resulting in some of the largest semileptonic B_s^0 yields. Fig. 1 shows the yield for the decay $B_s^0 \rightarrow D_s^- \mu^+ X (D_s^- \rightarrow \phi \pi^-)$ ¹ in $\sim 250 \text{ pb}^{-1}$ of data. The decay $B_s^0 \rightarrow D_s^- \mu^+ X (D_s^- \rightarrow K^{*0} K^-)$ has also been reconstructed and is expected to contribute significantly to the total B_s^0 yield. Other decays including hadronic B_s^0 decays are being studied as well.




Figure 1: $B_s^0 \rightarrow D_s^- \mu^+ X (D_s^- \rightarrow \phi \pi^-)$ yield.

¹Conjugate modes are implied throughout the paper

Other elements essential to a mixing analysis are given by the expression for the average statistical significance:

$$Significance(\Delta m_s, \sigma_t) = \sqrt{\frac{S\epsilon D^2}{2}} \sqrt{\frac{S}{S+B}} e^{-(\Delta m_s \sigma_t)^2/2} \quad (2)$$

where S is the number of signal events, ϵD^2 is a measure of the effectiveness of a flavor tagger (ϵ is the tagging efficiency while D is the “Dilution”), B is the number of background events and σ_t is the proper time resolution.

As indicated above, tagging the meson flavor (B_s^0 or $\bar{B}_s^0$) at decay time (final-state tagging) and at production time (initial-state tagging) are crucial. For the semileptonic modes used for mixing studies at DØ the final state particles provide the decay-time tag. For initial-state tagging, different techniques are being studied and optimized by doing measurements of Δm_d . Three tagging algorithms have been developed so far: opposite-side muon tagging, opposite-side jet charge tagging, and same-side (soft-pion) tagging.

The muon tagger relies on identifying the flavor of the other B meson in the event using the sign of the muon it decayed to - a negative muon corresponds to a b quark, and vice-versa. For the decay used for B_d^0 mixing studies - $B_d^0 \rightarrow D^{*-} \mu X$ where $D^{*-} \rightarrow \bar{D}^0 \pi^-$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ - both muons having the same sign would indicate that one B hadron had oscillated while opposite signs would indicate that neither (or both) had oscillated. Fig. 2 shows the measured asymmetry between the non-oscillated and oscillated mesons as a function of the visible proper decay length (VPDL). The fit to the asymmetry gives $\Delta m_d = 0.506 \pm 0.055 \pm 0.049 \text{ ps}^{-1}$ with an efficiency of $4.8 \pm 0.2 \%$ and $D = 46.0 \pm 4.2 \%$.




Figure 2: The asymmetry between the non-oscillated and oscillated mesons as a function of the visible proper decay length.

The opposite-side jet charge tagging algorithm calculates a p_T -weighted average charge of tracks: $\text{jetQ} \equiv \frac{\sum p_T^i q^i}{\sum p_T^i}$. A $|\text{jetQ}| < 0$ value corresponds to a b

quark for the B meson on the opposite side, and vice-versa. The same-side pion tagging algorithm makes use of the fact that the charge of the fragmentation pion correlates with the flavor of the reconstructed B meson. A positive pion corresponds to a $\bar{b}$ quark (i.e. B_d^0) if the reconstructed B meson is neutral, but to a b quark (i.e. B^-) if the reconstructed B meson is charged, and vice versa. Oscillations have been seen using both the opposite-side jet-charge and the same-side pion taggers, and work is ongoing to compute systematic errors and optimize performance. Lastly, since the significance scales as $e^{-(\Delta m_s \sigma_t)^2/2}$, for large values of Δm_s , a precise measurement of the proper decay time is crucial. Efforts are ongoing in this direction and we hope to have preliminary Δm_s results soon.

3 Conclusions

A preliminary measurement of Δm_d has been made with the upgraded DØ detector. The opposite-side soft muon tagger was used to obtain $\Delta m_d = 0.506 \pm 0.055 \pm 0.049$ ps⁻¹. This measurement, along with the development and optimization of other tagging algorithms, and the reconstruction of different $B_s \rightarrow D_s^\mp \mu^\pm X$ decay modes, is an important step towards a B_s mixing measurement.

References

1. D.Abbaneo, “Review of Experimental Results on Neutral B meson oscillations,” arXiv:hep-ex/0012010.
2. S.Eidelman *et al.*, “The Review of Particle Physics by the Particle Data Group,” Phys. Lett. **B592**, 1 (2004).
3. O.Schneider, “ $B^0 - \bar{B}^0$ mixing,” arXiv:hep-ex/0012010.